

Unclassi	ified						- 1
SECURITY CL	ASSIFICATION C	OF THIS PAGE					(
		REPORT	DOCUMENTATIO	N AC	)-A23	1 44	4. oved '04-0188
is REPORT ! Unclassi	SECURITY CLAS	SIFICATION		T1t	•		
2a. SECURITY	CLASSIFICATIO	N AUTHORITY			AVAILABILITY OF		
26. DECLASSI	FICATION / DOV	VNGRADING SCHEDU	ILE	unlimited	or buptic te	elease; d	istribution
4 PERFORMI	NG ORGANIZAT	TION REPORT NUMBE	ER(S)	5 MONITORING	ORGANIZATION RE	The same of	(5)
PL-TR-91	-2010					レロ	
Ceophys	PERFORMING ics Direc s Laborate	ORGANIZATION torate, ory.	6b OFFICE SYMBOL (If applicable) LID	73. NAME OF MONITORING AS WEATOF CITE FEB 0. 4' 1991			
	(City, State, an			76. ADDRESS (City, State, art. Zij, bde)			
	setts 017	31-5000		1		D	
85 NAMÉ OF ORGANIZ	FUNDING/SPC	ONSORING	36 OFFICE SYMBOL	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			
CRGAIVIZ.	AIICIA		(if applicable)				
BC ADDRESS	(City, State, and	d ZIP Code)	<u> </u>	10. SOURCE OF FUNDING NUMBERS			
				PROGRAM ELEMENT NO	NO NO	TASK NO	ACCESSION NO
				61102F	2310	G3	20
	lude Security C Reaction	15	otope Studies at Several Tem	and Energy D peratures	ependences	of Rate C	Constants
12 PERSONA	• •						
3a TYPE OF	REPORT	, A.A. Viggi	ano, John F. P Overed	14 DATE OF REPORT (Year, Month, Day) 15. PAGE COUNT			
Reprint		FROM	10	1991 Janua	ry 23		2
16 SUPPLEM	ENTARY NOTA	HON *Air For	ce Geophysics	Scholar -	Reprinted f	rom Noneq	uilibrium
Ettects	in Ion an	d Electron T	ransport, ed.	J.W. Gallagh	er et al.,	Plenem Pr	ess NY 1990
17	CUSATI	CODES	18. SUBJECT TERMS	(Continue on revers	e if necessary and	identify by blo	ock number)
FIELD	GROUP	SU8-GROUP	Ion-molecul	e reactions,	Isotope ex	change.	
	<u> </u>			ergy effects		0.	
19 ABSTHAC	T (Continue on	reverse it necessary	and identify by block i	lumber)			
			•				
	Abstract: F	Rate constant	s for the reactio	n O + N2O v	were measui	red using	a
1	emperatu	re variable se	lected ion flow	drift tube app	aratus. The	rate	
	•			• •			v
constants depend only upon total energy, not upon the form of that energy, i.e., whether internal or translational. The pure temperature dependence of							
the rate constants is $T^{-0.5}$ . The dominant product channel is $NO^- + NO$ .							
	ine rate CC	ו 15 מוונט וט ו	· THE COMMINA	ant product of	iannen 15 NO	, + NO.	

20 DISTRIBUTION/AVAILABILITY OF ABSTRACT  UNCLASSIFIED/UNLIMITED  SAME AS RPT.  OTIC USERS	21. ABSTRACT SECURITY CLASSIFICATION Unclassified
22a NAME OF RESPONSIBLE INDIVIDUAL	22b. TELEPHONE (Include Area Code) 22c. OFFICE SYMBOL
John F. Paulson	(617) 377-3124 LTD

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE
Unclassified

ISOTOPE STUDIES AND ENERGY DEPENDENCES OF RATE CONSTANTS FOR THE REACTION  $0^-$  +  $N_2O$  AT SEVERAL TEMPERATURES

Robert A. Morris\*, A. A. Viggiano and John F. Paulson Ionospheric Physics Division Geophysics Laboratory Hanscom AFB, MA 01731-5000

Rate constants for the reaction  $0^- + N_2 0$  were measured using a temperature variable-selected ion flow drift tube (SIFDT) instrument. The reaction was studied as a function of kinetic energy at four temperatures: 143, 196, 295, and 515 K. The product branching ratios of the reactions  $^{16}0^- + ^{14}N^{15}N^{16}0$  and  $^{18}0^- + ^{14}N^{15}N^{16}0$  were measured at two temperatures: 143 and 298 K.

The experiment employs a temperature varible flow tube with which to control the thermal energy of the reagents as well as an electric drift field to increase independently the kinetic energy of the ionic reactant. This permits fixing the overall center of mass collision energy while varying the relative contributions to that collision energy from the thermal energy of the reagents (due to temperature) and from the additional ion kinetic energy due to the electric field of the drift tube. Thus, reactions of monatomic ions (no internal modes) can be probed for the effects of the internal energy of the reactant neutral (Viggiano et al., 1988).

Figure 1 shows the rate constants for the reaction  $0^{-} + N_2 0$  plotted versus center-of-mass collision energy. The energy dependences of the rate constants measured at different experimental temperatures fall on the same curve within experimental uncertainty. Therefore, there appears to be little or no effect of the internal energy of the  $N_2 0$  reactant on the rate constant. In other words, the rate constant appears to depend only upon total energy and not on the particular type of energy. The pure temperature dependence of the rate constant is shown in the figure as a solid line and can be represented as  $T^{-0.5}$ .

Air Force Geophysics Scholar





Fig. 1. Rate constants for the reaction 0 + N<sub>2</sub>0 plotted as a function of center-of-mass kinetic energy.

The dominant product channel at the energies accessible in the experiment is that producing NO + NO. For the reaction of O with  $^{14}\mathrm{N}^{15}\mathrm{NO}$ , the ratio of the ionic products,  $^{14}\mathrm{N}^{0}/^{15}\mathrm{NO}$ , approached unity at zero  $^{14}\mathrm{N}^{15}\mathrm{NO}$  flow rate but decreased below unity with increasing  $^{14}\mathrm{N}^{15}\mathrm{NO}$  flow rate. It is suggested that the decrease is due to the secondary reaction  $^{14}\mathrm{N}^{0}$  +  $^{14}\mathrm{N}^{15}\mathrm{NO}$   $\longrightarrow$   $^{15}\mathrm{NO}$  +  $^{14}\mathrm{N}^{14}\mathrm{NO}$ , which we have measured to occur with a rate constant of 1 X 10  $^{-11}$  cm  $^3$  s  $^{-1}$ .

For the reaction  $^{18}0^- + ^{14}N^{15}N^{16}0$ , the 0 exchange product,  $^{16}0^-$ , was observable and represents a major product channel along with that producing NO + NO. All four NO isotopic products were observed:  $^{14}N^{16}0^-$ ,  $^{15}N^{16}0^-$ ,  $^{14}N^{18}0^-$ , and  $^{15}N^{18}0^-$ . The secondary reactions of both the exchange product  $^{16}0^-$  and the various NO products with N<sub>2</sub>O complicate the interpretation of the product spectra. Computer modeling of the coupled reactions and the resulting product branching ratios measured at the two experimental temperatures will be discussed.

## REFERENCES

Viggiano, A. A., R. A. Horris, and J. F. Paulson, 1988, J. Chem. Phys. 89, 4848; ibid. (in press, 1989).



400